

## Erratum

# Erratum to “The effects of nonnormality on asymptotic distributions of some likelihood ratio criteria for testing covariance structures under normal assumption” [J. Multivariate Anal. 96 (2005) 237–264]☆

Hirokazu Yanagihara<sup>a</sup>, Tetsuji Tonda<sup>b,\*</sup>, Chieko Matsumoto<sup>c,1</sup>

<sup>a</sup>Graduate School of Systems and Information Engineering, University of Tsukuba 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8573, Japan

<sup>b</sup>Department of Environmetrics and Biometrics, Research Institute for Radiation Biology and Medicine, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8553, Japan

<sup>c</sup>Department of Mathematics, Graduate School of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, Japan

The publisher regrets that the following corrections requested by the author at the proof stage were overlooked:

- Page 9, Line 19:

$$\frac{T_1 - (n-1)\eta_1}{\sqrt{n[\beta\tau_1^2 + 2(\beta-1)\{\text{tr}(\Sigma_0^{-1}\Sigma) - p\}^2]}} \rightarrow \frac{T_1 - (n-1)\eta_1}{\sqrt{n[\beta\tau_1^2 + (\beta-1)\{\text{tr}(\Sigma_0^{-1}\Sigma) - p\}^2]}}.$$

- Page 12, Line 19:

$$\frac{T_2 - (n-1)\eta_2}{\sqrt{n[\beta\tau_2^2 + 2(\beta-1)\{p\text{tr}(\Sigma)/\text{tr}(\Sigma) - p\}^2]}} \rightarrow \frac{T_2 - (n-1)\eta_2}{\sqrt{n\beta\tau_2^2}}.$$

- Page 13, Line 11:

$$\tau_3^2 = \text{tr}(\{\Sigma^{1/2}\Sigma_{(d)}^{-1}\Sigma^{1/2} - I_p\}^2) \rightarrow \text{tr}(\{\Sigma_{(d)}^{-1}\Sigma - I_p\}^2).$$

☆ DOI of original article: 10.1016/j.jmva.2004.10.014.

\* Corresponding author.

E-mail address: [ttetsuji@hiroshima-u.ac.jp](mailto:ttetsuji@hiroshima-u.ac.jp) (T. Tonda).

<sup>1</sup> Present address: Institute of Economic Research, Hitotsubashi University, 2-1 Naka, Kunitachi, Tokyo 186-8603, Japan.

- Page 16, Line 11:

$$\frac{T_3 - (n-1)\eta_3}{\sqrt{n[\beta\tau_3^2 + 2(\beta-1)\{\text{tr}(\mathbf{\Sigma}_{(d)}^{-1}\mathbf{\Sigma}) - p\}^2]}} \rightarrow \frac{T_3 - (n-1)\eta_3}{\sqrt{n\beta\tau_3}}.$$

- Page 17, Line 1:

$$\tau_4^2 = \sum_{i=1}^m \text{tr} \left( \rho_i^2 \{ \mathbf{\Sigma}_i^{1/2} \tilde{\mathbf{\Sigma}}^{-1} \mathbf{\Sigma}_i^{1/2} - \mathbf{I}_p \}^2 \right) \rightarrow \tau_4^2 = \sum_{i=1}^m \rho_i^2 \text{tr} \left( \{ \tilde{\mathbf{\Sigma}}^{-1} \mathbf{\Sigma}_i - \mathbf{I}_p \}^2 \right).$$

- Page 20, Line 7:

$$\begin{aligned} & \frac{T_4 - \eta_4}{\sqrt{n \left[ \beta\tau_4^2 + 2(\beta-1) \sum_{i=1}^m \left\{ \text{tr} \left( \mathbf{\Sigma}_i^{1/2} \tilde{\mathbf{\Sigma}}^{-1} \mathbf{\Sigma}_i^{1/2} - \mathbf{I}_p \right) \right\}^2 \right]}} \\ & \rightarrow \frac{T_4 - \eta_4}{\sqrt{n \left[ \beta\tau_4^2 + (\beta-1) \sum_{i=1}^m \rho_i^2 \left\{ \text{tr} \left( \tilde{\mathbf{\Sigma}}^{-1} \mathbf{\Sigma}_i \right) - p \right\}^2 \right]}}. \end{aligned}$$